

# MAU22302/33302 - Euclidean and non-Euclidean Geometry

## Homework 1

Trinity College Dublin

Course homepage Answers are due for February 11<sup>th</sup>, 23:59.

The use of electronic calculators and computer algebra software is allowed.

### **Exercise 1** *Playfair's Postulate (50pts)*

Euclid's formulation of the fifth postulate is rather lengthy:

*If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines - if produced indefinitely - meet on that side on which are the angles less than two right angles.*

It usually replaced by Playfair's postulate:


*Given a line  $\ell$  and a point  $P$  not on  $\ell$ , there exists exactly one line through  $P$  that does not intersect  $\ell$ , if both lines are extended indefinitely, i.e. there is a unique line parallel to  $\ell$  through  $P$ .*

The goal of this question is to establish their equivalence. You may freely use any of the postulates and any of propositions 1-27 given at the end of the assignment. In particular, you may assume at least one line parallel to  $\ell$  through a point  $P$  exists.

1. (20pts) Assume Postulates 1-5, and consider a line  $\ell$  and a point  $P$  not on  $\ell$ . By considering two lines through  $P$ , show that at least one intersects  $\ell$ . Conclude Playfair's Postulate.

*Hint: Drop a perpendicular from  $P$*

2. (20pts) Assume Postulates 1-4 and Playfair's Postulate. Establish that, if  $\ell_1$  and  $\ell_2$  are parallel, then  $\phi + \theta$  is equal to two right angles in the diagram below.



*Hint: Suppose otherwise and construct another line such that the corresponding angles add to two right angles*

3. (10pts) Hence conclude Postulate 5.

## Exercise 2 Triangles and Parallelograms (50pts)

In the following you may freely assume any of the given postulates and propositions, as well as your preferred parallel postulate from the first exercise

1. (10pts) Establish that the sum of angles in a triangle is equal to two right angles

*Hint: Introduce some alternate angles*

2. (10pts) Let  $\ell_1$  and  $\ell_2$  be parallel lines, with points  $A, B$  on  $\ell_1$  and  $C, D$  on  $\ell_2$ , appearing in that order. Suppose  $|AB| = |CD|$ . Show that  $AC$  is parallel to  $BD$ .

*Hint: BC27*

3. (15pts) Let  $\ell_1$  and  $\ell_2$  be parallel lines. Let  $A, B, E, F$  be points on  $\ell_1$ , appearing in that order. Let  $C, D$  be points on  $\ell_2$  in that order. Suppose

$$|AB| = |CD| = |EF|$$

Show that the parallelograms  $ABDC$  and  $EFDC$  have equal areas.

*Hint: Write each parallelogram as a sum/difference of triangles*

4. (15pts) Let  $\|\cdot\| : \mathbb{R}^n \rightarrow \mathbb{R}$  be a norm satisfying parallelogram rule

$$2\|x\|^2 + 2\|y\|^2 = \|x + y\|^2 + \|x - y\|^2$$

Show that the function

$$I(x, y) = \frac{1}{4} (\|x + y\|^2 - \|x - y\|^2)$$

is additive in the first variable, i.e.

$$I(x + y, z) = I(x, z) + I(y, z)$$

for all  $x, y, z \in \mathbb{R}^n$ .

*Not-a-hint: Draw a picture in  $\mathbb{R}^2$  to figure out why parallelogram rule is called that. It won't help with the problem, but it's nice to know!*

*Hint: You'll need to combine four instances of the parallelogram rule*

5. (0pts) Show that if  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  satisfies

$$f(x + y) = f(x) + f(y)$$

then  $f(qx) = qf(x)$  for all  $q \in \mathbb{Q}$ .

Suppose further that  $f$  is continuous. Show  $f(\lambda x) = \lambda f(x)$  for all  $\lambda \in \mathbb{R}$ . Hence conclude the function defined in the previous part is linear in the first variable.

# Postulates and propositions - slightly rephrased

## Postulates

1. There exists a line between any two points
2. A finite line segment can be extended indefinitely
3. Given a point and a line segment from that point, a circle with centre the given point and radius the given line segment can be drawn
4. All right angles are equal

## Propositions

1. One can construct an equilateral triangle on any base.
2. One can construct a line of length equal to a given line, from a given point
3. One can construct a line of length equal to the difference of two given line segments
4. If two triangles have two sides equal to two sides, and the angles contained are equal, then the triangles are equal
5. In an isosceles triangle, the angles at the base are equal
6. If two triangles have three sides equal to three sides, the triangles are equal
7. One can bisect a given angle
8. One can bisect a given finite line
9. One can draw a straight line at right angles to a given straight line from a given point on the line
10. One can draw a straight line at right angles to a given (infinite) straight line from a given point not on it.
11. If a straight line intersects a straight line, then it makes angles whose sum equals two right angles on the same side of a given line.

16. In angle triangle, if one of the sides is extended, the exterior angle is greater than either of the interior opposite angles
17. In any triangle, the sum of any two angles is less than two right angles.
23. One can construct an angle equal to a given angle on a given straight line and at a point on it.
27. If a straight line falling on two straight lines makes the alternate angles equal to one another, then the straight lines are parallel to one another
29. A straight line falling on two parallel straight lines makes the alternate angles equal to one another and the exterior angle equal to the interior opposite angle
31. Given a straight line and a point not on the line, one can draw a parallel line through the point