

MAU22103/33101 - Introduction to Number Theory

Exercise Sheet 1

Trinity College Dublin

Course homepage

Answers are due for October 1st, 16:00.

A physical copy should be submitted to the School of Mathematics office. If physical submission is not possible, a digital copy may be sent directly to me at akeilthytd.ie

The use of electronic calculators and computer algebra software is allowed.

Exercise 1 *Spare change*

Imagine a world in which we have exactly two types of coin: one worth P cents, one worth Q cents, such that $\gcd(P, Q) = 1$. The goal of this problem is to bound how many values we cannot form using these coins.

- (i) (15pts) Let us first consider $P = 5$, $Q = 13$. Determine in how many ways we can form 48 cents using these two coins

Hint: Consider the Diophantine equation $5u + 13v = 48$. What is the general form of the solution? We can't have a negative number of coins!

- (ii) (20pts) Show that we can form 49, 50, 51, 52 cents using coins worth $P = 5$ and $Q = 13$. Hence conclude that we can form any value $N \geq 48$.
- (iii) (15pts) Can we form 47 with coins worth $P = 5$ and $Q = 13$?

- (iv) (20pts) Define $g(P, Q) = PQ - P - Q$. Show that $g(P, Q)$ cannot be formed using coins worth P and Q
- (v) (20pts) Suppose that $P < Q$. Show that, for $N = 1, 2, \dots, P$, we can write

$$N = aP + bQ$$

with

$$0 < a < Q, \quad b + P > 0.$$

Hint: Start by showing we can find a solution with such an a and show that corresponding b satisfies the given inequality. Recall that $N - bQ > -bQ$

- (vi) (10pts) Hence conclude that $g(P, Q)$ is the largest value that cannot be formed using coins with P and Q .

This was the only exercise that you must submit before the deadline. The following exercises are not mandatory; they are not worth any points, and you do not have to submit them

However, I strongly encourage you to give them a try, as the best way to learn number theory is through practice.

The exercises marked with a star are the exercises I have suggested prioritising in the tutorials lecture. If there are any exercises you would particularly like to discuss, please let your TA

The exercises are arranged by theme, and roughly in order of difficulty within each theme, with the first few working as good warm-ups, and the remainder being of similar difficulty to the main exercise. You are welcome to email me if you have any questions about them. The solutions will be made available shortly after tutorials.

Exercise 2 Divisibility ★

Prove the following for $n \in \mathbb{N}$:

- (i) $5|n$ if and only if the last digit of n (in base 10) is 0 or 5.
- (ii) $2|n$ if and only if 2 divides the last digit (in base 10) of n .

- (iii) $4|n$ if and only if 4 divides the last two digits (in base 10) of n .
- (iv) $8|n$ if and only if 8 divides the last three digits (in base 10) of n .
- (v) $2^k|n$ if and only if 2^k divides the last k digits (in base 10) of n .
- (vi) $3|10a - 2$ if $3|a + 1$.
- (vii) A prime number k divides $n^2 - 1$ if and only if $k|n - 1$ or $k|n + 1$.

Exercise 3 *Applying the algorithm* ★

In the following, try to resolve the question without explicitly finding roots of the polynomials.

- (i) Prove that $4x^2 - 2x + 13$ has no integer roots.
- (ii) Prove that $3x^3 - 282x^2 + 18x - 28$ has no integer roots.
- (iii) Prove that, for any integers s, t, u, v , at least one of st , $sv + tu$, and uv is even. Hence conclude $ax^2 + bx + c$ has no rational roots if a, b, c are all odd.

Exercise 4 *Applying the algorithm* ★

Compute the following:

$$\begin{aligned}
 & i) \gcd(72, 18), \quad ii) \gcd(168, 124), \quad iii) \gcd(1047, 282), \\
 & iv) \gcd(n, 2n + 1), \quad v) \gcd(21n + 4, 14n + 3).
 \end{aligned}$$

Exercise 5 *Bézout's Theorem applied*

Prove that 2024 and 285 are coprime and determine integers u and v such that

$$2024u + 285v = 1.$$

Exercise 6 *Antimatter stamps are forbidden* ★

How many ways are there to pay for exactly €10.73 worth of postage if the post office will only sell you stamps worth 10c and 15c? How many ways are there to pay for exactly €10.75 worth of postage?

Note: The post office will not give change, even in the form of stamps. You cannot give antimatter stamps to cancel out regular stamps. You have to use a non-negative integer number of the stamps available to you.

Exercise 7 *Coprime products*

Let $a, b, c \in \mathbb{N}$. Prove that if a and b are coprime, and a and c are coprime, then a and bc are coprime

Exercise 8 *Primes and factorisations*

- (i) Write 2024 as a product of prime factors, and explain why each factor is prime.
- (ii) Hence or otherwise, work out the number of divisors of 2024 and their sum.
- (iii) Check the time on a 24 hour clock. Do the above computations for the time interpreted as a four digit number.
- (iv) Find all integers $M \in \mathbb{N}$ of the form $3^a 5^b$ such that the sum of the positive divisors of M is 33883.
Hint: $33883 = 31 \times 1093$, and both factors are prime.
- (v) Determine all $n \geq 2$ for which $n^2 - 1$ is prime.

Exercise 9 *Valuations of primes*

- (i) Let $m = \prod_i p_i^{a_i}$, $n = \prod_i p_i^{b_i}$ be two integers, where the p_i are pairwise distinct primes. Prove that $m \mid n$ iff. $a_i \leq b_i$ for each i .
Hint: If $n = km$, consider the prime factorisation of k .

- (ii) In what follows, let $p \in \mathbb{N}$ be prime. Recall that for nonzero $n \in \mathbb{Z}$, we define $v_p(n)$ as the exponent of p in n . Prove that for all nonzero $n \in \mathbb{Z}$, $v_p(n)$ is the largest integer v such that $p^v \mid n$.
- (iii) Recall that we set $v_p(0) = +\infty$ by convention. In view of the previous question, does this convention seem appropriate?
- (iv) Let $m, n \in \mathbb{Z}$, both nonzero. Prove that $v_p(mn) = v_p(m) + v_p(n)$. What happens if m or n is zero?
- (v) Let $m, n \in \mathbb{Z}$, both nonzero. Prove that $v_p(m+n) \geq \min(v_p(m), v_p(n))$. What happens if m or n is zero?
- (vi) Let $m, n \in \mathbb{Z}$. Prove that if $v_p(m) \neq v_p(n)$, then $v_p(m+n) = \min(v_p(m), v_p(n))$.
- (vii) Give an example where $v_p(m+n) > \min(v_p(m), v_p(n))$.

Exercise 10 Rational square roots ★

Prove that for $n \in \mathbb{N}$, either $\sqrt{n} \in \mathbb{N}$ or $\sqrt{n} \notin \mathbb{Q}$ is irrational.

Hint: First show that if $\sqrt{n} \notin \mathbb{N}$, then $v_p(n)$ is odd for some prime p . Then suppose $\sqrt{n} = \frac{a}{b}$ for some $a, b \in \mathbb{N}$. Rearrange this into an equality where you can compare valuations.

Exercise 11 Mersenne and perfect numbers

Define the Mersenne numbers by $M_0 = 0$, $M_1 = 1$, and

$$M_{n+1} = 3M_n - 2M_{n-1} \quad \text{for all } n \geq 1.$$

The Mersenne numbers are closely related to perfect numbers: n such that the sum of divisors of n is equal to $2n$: $\sigma_1(n) = 2n$. We will try to make this connection explicit.

1. Show that M_n is odd for all $n \geq 1$
2. Hence, show that $\gcd(M_n, M_{n+1}) = 1$ for every $n \in \mathbb{N}$.
3. Derive an explicit formula for M_n in terms of n .

Given an even natural number n , we can write it in the form $n = 2^a b$, where $a, b \in \mathbb{N}$, and b is odd.

4. Express $\sigma_1(n)$ in terms of a and $\sigma_1(b)$.
5. Hence show that if M_{a+1} is prime for some $a \geq 1$, the integer

$$n = M_{a+1}(M_a + 1)$$

is perfect.

We claim that every even perfect number is of this form.

6. Show that if $n = 2^a b$ is perfect, then $M_{a+1} | 2n$
7. Hence conclude $b = M_{a+1}c$ for some $c \in \mathbb{N}$. Show that $\sigma_1(b) = b + c$.
8. Explain why we must have $c = 1$, and $b = M_{a+1}$ is prime.
9. Hence determine an example of a perfect number $n > 6$

Exercise 12 Ideals of $\mathbb{Z}$

In this exercise, we define an *ideal* of $\mathbb{Z}$ to be a subset $I \subseteq \mathbb{Z}$ such that

- I is not empty,
 - whenever $i \in I$ and $j \in I$, we also have $i + j \in I$,
 - whenever $k \in \mathbb{Z}$ and $i \in I$, we also have $xi \in I$.
- (i) Let $n \in \mathbb{Z}$. Prove that $n\mathbb{Z} = \{nk, k \in \mathbb{Z}\}$ is an ideal of $\mathbb{Z}$.
 - (ii) For which $m, n \in \mathbb{Z}$ do we have $m\mathbb{Z} = n\mathbb{Z}$?
 - (iii) Let $I \subset \mathbb{Z}$ be an ideal. Prove that whenever $i \in I$ and $j \in I$, we also have $-i \in I$, $i - j \in I$, and $0 \in I$.
 - (iv) Let $I \subset \mathbb{Z}$ be an ideal. Prove that there exists $n \in \mathbb{Z}$ such that $I = n\mathbb{Z}$.
Hint: If $I \neq \{0\}$, let n be the smallest positive element of I , and consider the Euclidean division of the elements of i by n .
 - (v) Prove that if I and J are ideals of $\mathbb{Z}$, then

$$I + J = \{i + j \mid i \in I, j \in J\}$$

is also an ideal of $\mathbb{Z}$.

Hint: $i + j + i' + j' = i + i' + j + j'$.

- (vi) Let now $a, b \in \mathbb{Z}$. By the previous question, $a\mathbb{Z} + b\mathbb{Z}$ is an ideal, so it is of the form $c\mathbb{Z}$ for some $c \in \mathbb{Z}$. Express c in terms of a and b .

Hint: If you are lost, write an English sentence describing the set $a\mathbb{Z} + b\mathbb{Z}$.

- (vii) Prove that if I and J are ideals of $\mathbb{Z}$, then so is their intersection $I \cap J$.
- (viii) Let now $a, b \in \mathbb{Z}$. By the previous question, $a\mathbb{Z} \cap b\mathbb{Z}$ is an ideal, so it is of the form $c\mathbb{Z}$ for some $c \in \mathbb{Z}$. Express c in terms of a and b .