

MAU22103/33101 - Introduction to Number Theory

Exercise Sheet 1

Trinity College Dublin

Course homepage

Answers are due for October 1st, 16:00.

A physical copy should be submitted to the School of Mathematics office. If physical submission is not possible, a digital copy may be sent directly to me at akeilthytd.ie

The use of electronic calculators and computer algebra software is allowed.

Exercise 1 *Spare change*

Imagine a world in which we have exactly two types of coin: one worth P cents, one worth Q cents, such that $\gcd(P, Q) = 1$. The goal of this problem is to bound how many values we cannot form using these coins.

- (i) (15pts) Let us first consider $P = 5$, $Q = 13$. Determine in how many ways we can form 48 cents using these two coins

Hint: Consider the Diophantine equation $5u + 13v = 48$. What is the general form of the solution? We can't have a negative number of coins!

- (ii) (20pts) Show that we can form 49, 50, 51, 52 cents using coins worth $P = 5$ and $Q = 13$. Hence conclude that we can form any value $N \geq 48$.
- (iii) (15pts) Can we form 47 with coins worth $P = 5$ and $Q = 13$?

- (iv) (20pts) Define $g(P, Q) = PQ - P - Q$. Show that $g(P, Q)$ cannot be formed using coins worth P and Q
- (v) (20pts) Suppose that $P < Q$. Show that, for $N = 1, 2, \dots, P$, we can write

$$N = aP + bQ$$

with

$$0 < a < Q, \quad b + P > 0.$$

Hint: Start by showing we can find a solution with such an a and show that corresponding b satisfies the given inequality. Recall that $N - bQ > -bQ$

- (vi) (10pts) Hence conclude that $g(P, Q)$ is the largest value that cannot be formed using coins with P and Q .

Solution 1

- (i) We note that $\gcd(5, 13) = 1$ (they are both prime, and use Euclid's algorithm to find a solution to

$$5u + 13v = 1.$$

$$13 = 2(5) + 3$$

$$5 = 3 + 2$$

$$3 = 2 + 1$$

$$\Rightarrow$$

$$1 = 3 - 2$$

$$= 3 - (5 - 3) = 2(3) - 5$$

$$= 2(13 - 2 \times 5) - 5$$

$$= 2(13) - 5(5).$$

Multiplying by 48, we find

$$48 = 96(13) - 240(5)$$

Every other solution will be of the form

$$u = -240 + 13k, \quad v = 96 - 5k$$

for some k . In order to form 48 using coins, we need $u, v \geq 0$. Hence

$$18.4 \leq k \leq 19.2$$

and so $k = 19$ is the only solution.

$$u = 7, \quad v = 1.$$

- (ii) Since the question just asks if we can do this, it is enough to give any example of a non-negative solution in each case. The numbers are small enough that guessing works, but we present a more systematic method here.

Using the same method, we start with

$$98(13) - 245(5) = 49$$

$$100(13) - 250(5) = 50$$

$$102(13) - 255(5) = 51$$

$$104(13) - 260(5) = 52$$

and by adding $13k$ to the coefficient of 5 in each case so that we obtain a non-negative integer, we find

$$49 = 2(5) + 3(13)$$

$$50 = 10(5) + 0(13)$$

$$51 = 5(5) + 2(13)$$

$$52 = 0(5) + 4(13).$$

As we can form five consecutive values, 48, 49, 50, 51, 52, we can form any larger value by adding a multiple of 5 to one of these.

- (iii) We cannot! Starting from

$$(94)(13) + (-235)(5) = 47$$

we know that any integer solution must be of the form

$$u = 13k - 235, \quad v = 94 - 5k$$

If $u, v \geq 0$, we must have

$$k \geq 18.07, \quad k \leq 18.8.$$

However, no integers lie in this range, so no non-negative solutions can exist!

(iv) Suppose we can. Then we have $u, v \geq 0$ such that

$$PQ - P - Q = Pu + Qv$$

and hence

$$PQ = P(u + 1) + Q(v + 1).$$

This implies that

$$Q|P(u + 1) \text{ and } P|Q(v + 1).$$

As $\gcd(P, Q) = 1$, we must have

$$Q|(u + 1), \text{ and } P|(v + 1).$$

Thus, $PQ|P(u + 1), Q(v + 1)$. As $P(u + 1), Q(v + 1) > 0$, we must therefore have

$$P(u + 1), Q(v + 1) \geq PQ$$

but this is impossible. Therefore no such u, v exist.

(v) We know there exist integers a_0, b_0 such that

$$k = aP + bQ$$

and that, for any m

$$a = a_0 - Qm, \quad b = b_0 + Pm$$

is another solution. We can choose m so that $0 \leq a < Q$ (by division with remainder), and we cannot have $a = 0$ as $|bQ| \geq Q > P$ if b is also non-zero (or note that $Q \nmid k$).

We thus have

$$k - bQ = aP < PQ$$

Clearly $-bQ < k - bQ$, and so

$$-b < P \quad \Rightarrow \quad b + P > 0.$$

(vi) Hence, for any $k = 1, 2, \dots, P$, taking a and b as in the previous part we get

$$g(P, Q) + k = PQ - P - Q + aP + bQ = (a - 1)P + (b + P - 1)Q$$

gives the desired solution. For $N > P$, we can write $N = cP + k$ with $1 \leq k \leq P$ so that

$$g(P, Q) + N = (a + c - 1)P + (b + P - 1)Q$$

gives the necessary combination

This was the only exercise that you must submit before the deadline. The following exercises are not mandatory; they are not worth any points, and you do not have to submit them

However, I strongly encourage you to give them a try, as the best way to learn number theory is through practice.

The exercises marked with a star are the exercises I have suggested prioritising in the tutorials lecture. If there are any exercises you would particularly like to discuss, please let your TA

The exercises are arranged by theme, and roughly in order of difficulty within each theme, with the first few working as good warm-ups, and the remainder being of similar difficulty to the main exercise. You are welcome to email me if you have any questions about them. The solutions will be made available shortly after tutorials.

Exercise 2 *Divisibility* ★

Prove the following for $n \in \mathbb{N}$:

- (i) $5|n$ if and only if the last digit of n (in base 10) is 0 or 5.
- (ii) $2|n$ if and only if 2 divides the last digit (in base 10) of n .
- (iii) $4|n$ if and only if 4 divides the last two digits (in base 10) of n .
- (iv) $8|n$ if and only if 8 divides the last three digits (in base 10) of n .
- (v) $2^k|n$ if and only if 2^k divides the last k digits (in base 10) of n .
- (vi) $3|10a - 2$ if $3|a + 1$.
- (vii) A prime number k divides $n^2 - 1$ if and only if $k|n - 1$ or $k|n + 1$.

Solution 2

- (i) Let ℓ be the last digit of n . Then we can write $n = 10m + \ell$ for some integer $m \in \mathbb{Z}$. As $5|10m$, we must have that $5|n = 10m + \ell$ if and only if $5|\ell = n - 10m$. The only one digit numbers divisible by 5 are 0 and 5.
- (ii) Similarly to the last part, $2|n$ if and only if $2|\ell$ as $2|10m$.
- (iii) Let ℓ be integer given by the last two digits of n , so that we can write $n = 100m + \ell$ for some integer m . As $4|100$, $4|100m$, and hence $4|n$ if and only if $4|\ell$.
- (iv) Note that $8|1000$, and make the same argument.
- (v) If ℓ denotes the integer given by the last k digits of n , then we can write $n = 10^k + \ell$. Since $2^k|10^k$, we can apply the same argument to conclude that $2^k|n$ if and only if $2^k|\ell$.
- (vi) Note that $10a - 2 = 10(a + 1) - 12$. If $3|a + 1$, we must have $3|10a - 2$, as $3|-12$.
- (vii) We can factorise $n^2 - 1 = (n - 1)(n + 1)$. Clearly, if $k|n - 1$ or $k|n + 1$, then $k|n^2 - 1$. Since k is prime, if $k|(n - 1)(n + 1)$, then k divides at least one of the two factors.

Exercise 3 *Tricks for surviving the cube* ★

In the following, try to resolve the question without finding explicitly finding roots of the polynomials.

- (i) Prove that $4x^2 - 2x + 13$ has no integer roots.
- (ii) Prove that $3x^3 - 282x^2 + 18x - 28$ has no integer roots.
- (iii) Prove that, for any integers s, t, u, v , at least one of st , $sv + tu$, and uv is even. Hence conclude $ax^2 + bx + c$ has no rational roots if a, b, c are all odd.

Solution 3

- (i) Suppose $4n^2 - 2n + 13 = 0$ for some $n \in \mathbb{Z}$. Then $13 = 2(n - 2n^2)$, but $2 \nmid 13$. Therefore no such n exists.
- (ii) Note that $3|3$, $3| -282$, and $3|18$, but $3 \nmid -28$, and so by the same reasoning as the previous question, no integer roots can exist.
- (iii) Suppose all of st , $sv + tu$ and uv are odd. Then, since $2 \nmid st$ and $2 \nmid uv$, we must have that s, t, u, v are all odd. Therefore sv and tu are odd, and the sum of two odd numbers is even, so we must have $2|sv + tu$.
If $ax^2 + bx + c$ has a rational root, it can be factorised as $(sx + u)(tx + v)$ for some integers s, t, u, v such that $a = st$, $b = sv + tu$, $c = uv$. As noted above, if a and c are odd, b must be even. Therefore no such factorisation can exist.

Exercise 4 *Applying the algorithm* ★

Compute the following:

$$\begin{aligned} & i) \gcd(72, 18), \quad ii) \gcd(168, 124), \quad iii) \gcd(1047, 282), \\ & iv) \gcd(n, 2n + 1), \quad v) \gcd(21n + 4, 14n + 3). \end{aligned}$$

Solution 4

- (i) $72 = 4(18)$, so $\gcd(72, 18) = 18$.
- (ii) $168 = 124 + 44$, $124 = 2(44) + 36$, $44 = 36 + 8$, $36 = 4(8) + 4$, $8 = 2(4)$, so $\gcd(168, 124) = 4$.
- (iii) $1047 = 3(282) + 201$, $282 = 201 + 81$, $201 = 2(81) + 39$, $81 = 2(39) + 3$, $39 = 13(3)$, so $\gcd(1047, 282) = 3$.
- (iv) $2n + 1 = 2(n) + 1$, so $\gcd(2n + 1, n) = 1$.
- (v) $21n + 4 = 14n + 3 + 7n + 1$, $14n + 3 = 2(7n + 1) + 1$, so $\gcd(21n + 4, 14n + 3) = 1$.

Exercise 5 *Bézout's Theorem applied*

Prove that 2024 and 285 are coprime and determine integers u and v such that

$$2024u + 285v = 1.$$

Solution 5

We will apply Euclid's algorithm:

$$2024 = 7(285) + 29$$

$$285 = 9(29) + 24$$

$$29 = 24 + 5$$

$$24 = 4(5) + 4$$

$$5 = 4 + 1$$

and hence $\gcd(2024, 285) = 1$. To determine u and v , we run the algorithm backwards

$$\begin{aligned} 1 &= 5 - 4 \\ &= 5 - (24 - 4(5)) = 5(5) - 24 \\ &= 5(29 - 24) - 24 = 5(29) - 6(24) \\ &= 5(29) - 6(285 - 9(29)) = 59(29) - 6(285) \\ &= 59(2024 - 7(285)) - 6(285) = 59(2024) - 419(285). \end{aligned}$$

Thus, a solution is given by $(u, v) = (59, -419)$.

Exercise 6 *Antimatter stamps are forbidden* ★

How many ways are there to pay for exactly €10.73 worth of postage if the post office will only sell you stamps worth 10c and 15c? How many ways are there to pay for exactly €10.75 worth of postage?

Note: The post office will not give change, even in the form of stamps. You cannot give antimatter stamps to cancel out regular stamps. You have to use a non-negative integer number of the stamps available to you.

Solution 6

The first part of the question is asking us to find the number of non-negative integer solutions to the Diophantine equation

$$10u + 15v = 1073.$$

Note that $\gcd(10, 15) = 5 \nmid 1073$, so there are no integer solutions, let alone non-negative ones.

The second part of the question is asking us to find the number of non-negative integer solutions to the Diophantine equation

$$10u + 15v = 1075.$$

As $5 \mid 1075$, we know that infinitely many solutions exist. We can, in fact, reduce the problem to finding non-negative integer solutions to

$$2u + 3v = 215.$$

We will first find an integer solution to

$$2x + 3y = 1$$

which we could solve via the Euclidean algorithm, as in the proof of Bézout's theorem, but there is an obvious solution $(x, y) = (-1, 1)$. Hence, we can take $(u_0, v_0) = (-215, 215)$ as a solution to our main equation. Every other solution is of the form

$$(u, v) = (-215 + 3k, 215 - 2k)$$

as k ranges across $\mathbb{Z}$. In order to obtain a non-negative integer solution, we need that

$$3k \geq 215 \quad \text{and} \quad 215 \geq 2k$$

or equivalently that $107.5 \geq k \geq 71.6666$. Since k must be an integer, we can restrict to $107 \geq k \geq 72$. Every such k gives a non-negative integer solution, and every non-negative integer solution corresponds to such a k , so we must therefore have $107 - 71 = 36$ integer solutions. The easiest solution to count probably corresponds to $k = 107$, giving $(u, v) = (106, 1)$. The solution involving the least number of stamps corresponds to $k = 72$, with $(u, v) = (1, 71)$. Personally, I think $k = 86$, $(u, v) = (43, 43)$ is fun, even if everyone involved in the postage probably disagrees.

Exercise 7 *Coprime products*

Let $a, b, c \in \mathbb{N}$. Prove that if a and b are coprime, and a and c are coprime, then a and bc are coprime

Solution 7

Suppose $d|a$. Then, as $\gcd(d, b)|d$, we must have that $\gcd(d, b)|a$. By definition $\gcd(d, b)|b$, and so $\gcd(d, b)|\gcd(a, b) = 1$. Hence $\gcd(d, b) = 1$. Now, if $d|a$ and $d|bc$, then by Gauss' Lemma, $d|c$, as $\gcd(d, b) = 1$. But if $d|a$ and $d|c$, then $d|\gcd(a, c) = 1$ and so $d = 1$. Therefore a and bc must be coprime, as 1 is their only common divisor.

Exercise 8 *Primes and factorisations*

- (i) Write 2024 as a product of prime factors, and explain why each factor is prime.
- (ii) Hence or otherwise, work out the number of divisors of 2024 and their sum.
- (iii) Check the time on a 24 hour clock. Do the above computations for the time interpreted as a four digit number.
- (iv) Find all integers $M \in \mathbb{N}$ of the form $3^a 5^b$ such that the sum of the positive divisors of M is 33883.

Hint: $33883 = 31 \times 1093$, and both factors are prime.

- (v) Determine all $n \geq 2$ for which $n^2 - 1$ is prime.

Solution 8

- (i) We begin by noting that $2024 = 8(253)$: Exercise 2(iv) lets us deduce this quickly. Thus $2024 = 2^3 \cdot 253$. 2 is prime, as there aren't any smaller integers to be non-trivial factors.

To factor 253, we begin dividing by small primes less than or equal to $\sqrt{253} \approx 16$, and find that it is only divisible by 11. Thus

$$2024 = 2^3 \cdot 11 \cdot 23.$$

It is easy to check that 11 is prime, and 23 must be prime, either by direct computation or by noting that 253 would have to have a factor smaller than 11 if 23 were composite.

- (ii) Every divisor of 2024 is of the form $2^a \cdot 11^b \cdot 23^c$, where $0 \leq a \leq 3$, and $0 \leq b, c \leq 1$. This gives us $4 \times 2 \times 2 = 16$ divisors. Their sum is given by the product

$$\left(\frac{2^4 - 1}{2 - 1}\right) \left(\frac{11^2 - 1}{11 - 1}\right) \left(\frac{23^2 - 1}{23 - 1}\right) = 4320.$$

- (iii) It is currently $15 : 36$. Dividing repeatedly by 2, we find $1536 = 2^9 \cdot 3$. This means that 1536 has $10 \cdot 2 = 20$ divisors, and the sum of these divisors is given by

$$\frac{(2^{10} - 1)(9 - 1)}{2} = 4092.$$

- (iv) From our formula, the sum of divisors of $3^a 5^b$ is given by

$$\left(\frac{3^{a+1} - 1}{2}\right) \left(\frac{5^{b+1} - 1}{4}\right) = (1 + 3 + \cdots + 3^a) (1 + 5 + \cdots + 5^b)$$

Therefore if $\sigma_1(3^a 5^b) = 33883 = 31$, we must have that one of four cases occurs: both prime factors divide the 3-sum, both prime factors divide the 5 sum, 31 divides the 3-sum and 1093 divides the 5-sum, or 31 divides the 5-sum and 1093 divides the 3 sum.

In fact, as there are no other prime factors, we can replace “divides” with “is equal to”.

In the first case $b = 0$, and we need

$$1 + 3 + \cdots + 3^a = \frac{3^{a+1} - 1}{2} = 33883$$

and so $3^{a+1} = 2(33883) + 1 = 67767$, which is not a power of three.

In the second case $a = 0$, and we need

$$\frac{5^{b+1} - 1}{4} = 33883$$

and so $5^{b+1} = 4(33883) + 1 = 135533$, which is not even a multiple of 5.

In the third case, we must have

$$\frac{3^{a+1} - 1}{2} = 31 \quad \text{and} \quad \frac{5^{b+1} - 1}{4} = 1093$$

and hence $3^{a+1} = 63$, which is not a power of three.

In the final case, we must have

$$3^{a+1} = 2187 = 3^7 \quad \text{and} \quad 5^{b+1} = 125 = 5^3$$

Hence, $a = 6$ and $b = 2$ is the only possible solution, giving

$$M = 18225.$$

- (v) We have that $n^2 - 1 = (n - 1)(n + 1)$ is a factorisation of $n^2 - 1$, into positive integers, since $n \geq 2$. In order for $n^2 - 1$ to be prime, these two factors must be 1 and $n^2 - 1$. In particular, $n - 1 = 1$ and $n + 1 = n^2 - 1$. This gives $n = 2$, and so 3 is the only prime of the form $n^2 - 1$.

Exercise 9 *Valuations of primes*

- (i) Let $m = \prod_i p_i^{a_i}$, $n = \prod_i p_i^{b_i}$ be two integers, where the p_i are pairwise distinct primes. Prove that $m \mid n$ iff. $a_i \leq b_i$ for each i .
Hint: If $n = km$, consider the prime factorisation of k .
- (ii) In what follows, let $p \in \mathbb{N}$ be prime. Recall that for nonzero $n \in \mathbb{Z}$, we define $v_p(n)$ as the exponent of p in n . Prove that for all nonzero $n \in \mathbb{Z}$, $v_p(n)$ is the largest integer v such that $p^v \mid n$.
- (iii) Recall that we set $v_p(0) = +\infty$ by convention. In view of the previous question, does this convention seem appropriate?
- (iv) Let $m, n \in \mathbb{Z}$, both nonzero. Prove that $v_p(mn) = v_p(m) + v_p(n)$. What happens if m or n is zero?
- (v) Let $m, n \in \mathbb{Z}$, both nonzero. Prove that $v_p(m+n) \geq \min(v_p(m), v_p(n))$. What happens if m or n is zero?

- (vi) Let $m, n \in \mathbb{Z}$. Prove that if $v_p(m) \neq v_p(n)$, then $v_p(m + n) = \min(v_p(m), v_p(n))$.
- (vii) Give an example where $v_p(m + n) > \min(v_p(m), v_p(n))$.

Solution 9

- (i) Suppose $m|n$ so that $n = mk$. We can write $k = \prod_i p_i^{c_i}$ for some non-negative integers c_i , using the same set of primes, as if $p|k$ then $p|n$. Thus

$$\prod_i p_i^{b_i} = \left(\prod_i p_i^{a_i} \right) \left(\prod_i p_i^{c_i} \right) = \prod_i p_i^{a_i + c_i}$$

By the uniqueness of prime factorisation, we must have $a_i + c_i = b_i$ for each i , and since $c_i \geq 0$, we must have $a_i = b_i - c_i \leq b_i$.

Conversely, if $b_i \geq a_i$, then $c_i = b_i - a_i \geq 0$, and so

$$k = \prod_i p_i^{c_i}$$

is an integer such that $n = km$. Therefore $m|n$.

- (ii) By the previous part of the question, $p^v|n$ if and only if $v \leq v_p(n)$, from which the conclusion is immediate.
- (iii) Yes! There is no upper bound on integers v such that $p^v|0$, so declaring the valuation of 0 to be infinite works with this intuition.
- (iv) By the definition of $v_p(m)$ we can write $m = p^{v_p(m)}M$ where $p \nmid M$, and similarly for n . Then $mn = p^{v_p(m)+v_p(n)}MN$. Since $p \nmid M$ and $p \nmid N$, $p \nmid MN$ and hence $v = v_p(m) + v_p(n)$ is the maximal integer such that $p^v|mn$, which is to say $v_p(mn) = v_p(m) + v_p(n)$.

This extends to the case where m or n is zero (without loss of generality $m = 0$), as then $mn = 0$, so $v_p(mn) = \infty = \infty + v_p(n)$.

- (v) As before, write $m = p^{v_p(m)}M$ and $n = p^{v_p(n)}N$, and without loss of generality assume that $v_p(m) \leq v_p(n)$. We write $v_p(n) = v_p(m) + c$ for some $c \geq 0$. Then

$$m + n = p^{v_p(m)}M + p^{v_p(m)+c}N = p^{v_p(m)}(M + p^cN)$$

and so $p^{v_p(m)} \mid m+n$. As $v_p(m+n)$ is the maximal integer v such that $p^v \mid m+n$, we must have $v_p(m+n) \geq v_p(m) = \min(v_p(m), v_p(n))$.

- (vi) Take $m = n = 2$. Then $v_2(m) = v_2(n) = 1$, but $v_2(m+n) = 2$. In general, equality will hold if and only if $v_p(m) \neq v_p(n)$.

Exercise 10 *Rational square roots* ★

Prove that for $n \in \mathbb{N}$, either $\sqrt{n} \in \mathbb{N}$ or $\sqrt{n} \notin \mathbb{Q}$ is irrational.

Hint: First show that if $\sqrt{n} \notin \mathbb{N}$, then $v_p(n)$ is odd for some prime p . Then suppose $\sqrt{n} = \frac{a}{b}$ for some $a, b \in \mathbb{N}$. Rearrange this into an equality where you can compare valuations.

Solution 10

Note that $v_p(k^2) = v_p(k) + v_p(k) = 2v_p(k)$ is even, and hence the p -adic valuation of a square number is even for any prime p . Thus, if n is not a square, which is to say $\sqrt{n} \notin \mathbb{N}$, then $v_p(n)$ must be odd for some prime p .

Suppose $\sqrt{n} = \frac{a}{b}$. This implies that

$$a^2 = nb^2.$$

Let p be a prime such that $v_p(n)$ is odd. Then we must have

$$2v_p(a) = v_p(a^2) = v_p(nb^2) = v_p(n) + v_p(b^2) = v_p(n) + 2v_p(b).$$

But this implies $v_p(n)$ must be divisible by 2, a contradiction. Therefore $\sqrt{n}$ is irrational if it is not an integer.

Exercise 11 *Mersenne and perfect numbers*

Define the Mersenne numbers by $M_0 = 0$, $M_1 = 1$, and

$$M_{n+1} = 3M_n - 2M_{n-1} \quad \text{for all } n \geq 1.$$

The Mersenne numbers are closely related to perfect numbers: n such that the sum of divisors of n is equal to $2n$: $\sigma_1(n) = 2n$. We will try to make this connection explicit.

1. Show that M_n is odd for all $n \geq 1$

2. Hence, show that $\gcd(M_n, M_{n+1}) = 1$ for every $n \in \mathbb{N}$.
3. Derive an explicit formula for M_n in terms of n .
Given an even natural number n , we can write it in the form $n = 2^a b$, where $a, b \in \mathbb{N}$, and b is odd.
4. Express $\sigma_1(n)$ in terms of a and $\sigma_1(b)$.
5. Hence show that if M_{a+1} is prime for some $a \geq 1$, the integer

$$n = M_{a+1}(M_a + 1)$$

is perfect.

We claim that every even perfect number is of this form.

6. Show that if $n = 2^a b$ is perfect, then $M_{a+1} | 2n$
7. Hence conclude $b = M_{a+1}c$ for some $c \in \mathbb{N}$. Show that $\sigma_1(b) = b + c$.
8. Explain why we must have $c = 1$, and $b = M_{a+1}$ is prime.
9. Hence determine an example of a perfect number $n > 6$

Solution 11

1. Since $M_{n+1} = 3M_n - 2M_{n-1}$, M_{n+1} is divisible by 2 if and only if M_n is. Since $M_1 = 1$, this implies M_n is odd for all $n \geq 1$.
2. Since $M_{n+1} = 3M_n - 2M_{n-1}$, we must have that $\gcd(M_n, M_{n+1}) | 2M_{n-1}$. As M_{n+1} is odd for all $n \geq 0$, this implies $\gcd(M_n, M_{n+1}) | M_{n-1}$. Hence $\gcd(M_n, M_{n+1}) = \gcd(M_n, M_{n-1})$. Iterating this argument, we find that

$$\begin{aligned} \gcd(M_{n+1}, M_n) &= \gcd(M_n, M_{n-1}) = \gcd(M_{n-1}, M_{n-2}) \\ &= \cdots = \gcd(M_2, M_1) = \gcd(M_1, M_0) = 1. \end{aligned}$$

3. Looking at the first few values of M_n , we guess that $M_n = 2^n - 1$. Note that

$$3(2^n - 1) - 2(2^{n-1} - 1) = 3 \cdot 2^n - 3 - 2^n + 2 = 2^{n+1} - 1$$

so $2^n - 1$ satisfies the same recursion. As $M_n = 2^n - 1$ for $n = 0, 1$, we therefore have $M_n = 2^n - 1$ for all $n \geq 0$.

4. As b is odd, $\gcd(2^a, b) = 1$ and so $\sigma_1(2^a b) = \sigma_1(2^a) + \sigma_1(b)$. The prime power 2^a has $\sigma_1(2^a) = \frac{2^{a+1}-1}{2-1}$, and so

$$\sigma_1(n) = (2^{a+1} - 1)\sigma_1(b).$$

5. Knowing $M_n = 2^n - 1$, we are interested in the integer

$$n = 2^a(2^{a+1} - 1)$$

which is of the form $n = 2^a b$ for $b = 2^{a+1} - 1$ an odd integer. Thus

$$\sigma_1(n) = (2^{a+1}-1)\sigma_1(2^{a+1}-1) = (2^{a+1}-1)(2^{a+1}-1+1) = 2^{a+1}(2^{a+1}-1) = 2n$$

as $2^{a+1} - 1$ is prime.

6. Suppose n is perfect. Then

$$2n = \sigma(n) = (2^{a+1} - 1)\sigma_1(b) = M_{a+1}\sigma_1(b)$$

and so $M_{a+1} | 2n$.

7. By the previous part $M_{a+1} | 2^{a+1}b$. Since M_{a+1} is odd, and so $\gcd(M_{a+1}, 2^{a+1}) = 1$, we must have $M_{a+1} | b$. Thus, $b = M_{a+1}c$ for some $c \in \mathbb{N}$. Hence

$$M_{a+1}\sigma_1(b) = 2n = 2^{a+1}b = (M_{a+1} + 1)b = M_{a+1}b + b = M_{a+1}b + M_{a+1}c$$

and so $\sigma_1(b) = b + c$.

8. If $c > 1$, then $\sigma_1(b) \geq b + c + 1 > b + c$ as $c | b$. Thus $c = 1$ and $b = M_{a+1}$. If b is not prime, then it has a proper divisor $d > 1$ and so $\sigma_1(b) \geq b + d + 1 > b + 1 = b + c$. Thus, b must be prime.

9. Take $a = 2$. As $M_3 = 7$ is prime, $n = 2^2 \times 7 = 28$ must be perfect.

Exercise 12 Ideals of $\mathbb{Z}$

In this exercise, we define an *ideal* of $\mathbb{Z}$ to be a subset $I \subseteq \mathbb{Z}$ such that

- I is not empty,
- whenever $i \in I$ and $j \in I$, we also have $i + j \in I$,

- whenever $k \in \mathbb{Z}$ and $i \in I$, we also have $xi \in I$.
- (i) Let $n \in \mathbb{Z}$. Prove that $n\mathbb{Z} = \{nk, k \in \mathbb{Z}\}$ is an ideal of $\mathbb{Z}$.
- (ii) For which $m, n \in \mathbb{Z}$ do we have $m\mathbb{Z} = n\mathbb{Z}$?
- (iii) Let $I \subset \mathbb{Z}$ be an ideal. Prove that whenever $i \in I$ and $j \in I$, we also have $-i \in I$, $i - j \in I$, and $0 \in I$.
- (iv) Let $I \subset \mathbb{Z}$ be an ideal. Prove that there exists $n \in \mathbb{Z}$ such that $I = n\mathbb{Z}$.
Hint: If $I \neq \{0\}$, let n be the smallest positive element of I , and consider the Euclidean division of the elements of I by n .
- (v) Prove that if I and J are ideals of $\mathbb{Z}$, then

$$I + J = \{i + j \mid i \in I, j \in J\}$$

is also an ideal of $\mathbb{Z}$.

Hint: $i + j + i' + j' = i + i' + j + j'$.

- (vi) Let now $a, b \in \mathbb{Z}$. By the previous question, $a\mathbb{Z} + b\mathbb{Z}$ is an ideal, so it is of the form $c\mathbb{Z}$ for some $c \in \mathbb{Z}$. Express c in terms of a and b .
Hint: If you are lost, write an English sentence describing the set $a\mathbb{Z} + b\mathbb{Z}$.
- (vii) Prove that if I and J are ideals of $\mathbb{Z}$, then so is their intersection $I \cap J$.
- (viii) Let now $a, b \in \mathbb{Z}$. By the previous question, $a\mathbb{Z} \cap b\mathbb{Z}$ is an ideal, so it is of the form $c\mathbb{Z}$ for some $c \in \mathbb{Z}$. Express c in terms of a and b .

Solution 12

- (i) Clearly $n \in n\mathbb{Z}$, so $n\mathbb{Z}$ is non-empty. If $a, b \in n\mathbb{Z}$, then there exist $k, \ell \in \mathbb{Z}$ such that $a = kn$ and $b = \ell n$. Hence $a + b = n(k + \ell) \in n\mathbb{Z}$. If $a = nk \in n\mathbb{Z}$ and $m \in \mathbb{Z}$, then $ma = n(mk) \in n\mathbb{Z}$. Hence, $n\mathbb{Z}$ is an ideal of $\mathbb{Z}$.
- (ii) If $m\mathbb{Z} = n\mathbb{Z}$, then $m \in n\mathbb{Z}$, and so $n|m$. Similarly, we must have $m|n$. But if $m|n$ and $n|m$, we must have $m = \pm n$.

- (iii) If $i \in I$, then $(-1)i = -i \in I$, via the multiplicative property of ideals. If $j \in I$, then $-j \in I$, and since ideals are closed under addition $i + (-j) = i - j \in I$. Finally, since I is non-empty, there exists an element $k \in I$, and therefore $0 = 0(k) \in I$.
- (iv) If $I = \{0\}$, then $I = 0\mathbb{Z}$, so assume $I \neq \{0\}$. Then, I contains a smallest positive element n , and all multiple of n . Hence $n\mathbb{Z} \subset I$. Now suppose $N \in I$, which we can assume without loss of generality to be positive. Dividing N by n , we get $N = nq + r$, $0 \leq r < n$. Writing $r = N - nq$, we see that $r \in I$ as $N \in I$ and $nq \in I$. But if $r \neq 0$, this contracts the minimality of n . Hence $r = 0$, and $N = nq \in n\mathbb{Z}$. Thus $I \subset n\mathbb{Z}$ and $I = n\mathbb{Z}$.
- (v) Since $0 \in I$ and $0 \in J$, $0 \in I + J$, so $I + J$ is non empty. To see that it is closed under addition, suppose $k, \ell \in I + J$. Then there exists $i, i' \in I$ and $j, j' \in J$ such that $k = i + j$ and $\ell = i' + j'$. Hence

$$k + \ell = i + j + i' + j' = (i + i') + (j + j') \in I + J$$

as $(i + i') \in I$ and $(j + j') \in J$. Finally we check the multiplicative property. Suppose $k \in I + J$ and $m \in \mathbb{Z}$. Then $k = i + j$ for some $i \in I$ and $j \in J$, and hence $mk = mi + mj \in I + J$ as $mi \in I$ and $mj \in J$.

- (vi) The ideal $a\mathbb{Z} + b\mathbb{Z}$ given as a set is

$$\{au + bv \mid u, v \in \mathbb{Z}\}$$

which by (corollaries to) Bézout's theorem is precisely the set of multiples of $\gcd(a, b)$. Hence $c = \pm \gcd(a, b)$.

- (vii) Since $0 \in I$ and $0 \in J$, $0 \in I \cap J$, so it is non-empty. If $i, j \in I \cap J$, then $i + j \in I$ and $i + j \in J$, and so $i + j \in I \cap J$. If $i \in I \cap J$ and $k \in \mathbb{Z}$, then $ki \in I$ and $ki \in J$, so $ki \in I \cap J$.
- (viii) $a\mathbb{Z}$ is the set of integers divisible by a . $b\mathbb{Z}$ is the set of integers divisible by b . Their intersection is the set of integers divisible by both a and b . Hence $\text{lcm}(a, b) \in a\mathbb{Z} \cap b\mathbb{Z}$. Furthermore, every such integer is divisible by $\text{lcm}(a, b)$, and so $c = \pm \text{lcm}(a, b)$.