

MAU11S02 sixth Friday quiz, week 9

Friday 25/3/22 due 4pm Monday 28/3/22

Rules and procedures.

1. Attempt 3 questions. Only *your first three answers* will be marked. **2.** Each question carries 20 marks, so the maximum quiz mark is 60. **3.** If a particular method of solution is stipulated, you get no marks if you don't use it. **4. *Show all work.*** No marks will be given for answers which do not show the calculations. **5.** Your answers should be scanned and submitted to Blackboard as a 'Friday assignment.'

Question 1. Calculate the quadratic function $y = ax^2 + bx + c$ best fitting the following data (least squared error estimate). ***You must use the formula given in lectures, and show the calculations.***

$$(-3, 1) \quad (-2, 0) \quad (-1, 1) \quad (1, 3)$$

Question 2. Find eigenvalues and eigenvectors, and the change-of-basis matrix S , for the matrix A , where

$$A = \begin{bmatrix} 1 & 2 \\ 5 & -2 \end{bmatrix}$$

Question 3. Calculate e^{At} , where A is as in Question 2.

Question 4. Solve in full the differential equations

$$\frac{dx}{dt} = x + 2y; \quad \frac{dy}{dt} = 5x - 2y$$

with $x = 1$ and $y = 2$ at $t = 0$.

Question 5. There are 4 balls in one jar, labelled 2, 3, 4 respectively, and 3 balls in another jar, labelled 2, 7, 7, 8 respectively.

The random variables X_1, X_2 represent the labels x_1, x_2 on two balls chosen randomly from each jar.

(i) List the *nine* possible outcomes (x_1, x_2) and their probabilities.

(ii) List the *seven* possible values of $x_1 + x_2$ and their probabilities.

Question 6. The least-squared-error method can be used to find best-fitting curves of many different shapes. Find the best-fitting curve of the form $y = 2^x a + (-1)^x b$ for the data points in Question 1.

Show that $((A^T A)_{1,1})$ was given wrongly as 2.578125)

$$A^T A = \begin{bmatrix} 4.328125 & -2.375 \\ -2.375 & 4 \end{bmatrix}, \quad (\text{corrected}) \quad \text{and} \quad A^T Y = \begin{bmatrix} 6.625 \\ -5 \end{bmatrix}$$

and hence solve the problem.