

UNIVERSITY OF DUBLIN

XMA2161

TRINITY COLLEGE

FACULTY OF SCIENCE

SCHOOL OF MATHEMATICS

SF Mathematics
SF Theoretical Physics

Trinity Term 2007

COURSE 216

Thursday, May 24

Luce Hall

14.00 — 16.00

Dr. J. Stalker

ATTEMPT FOUR QUESTIONS

Log tables are available from the invigilators, if required.

Non-programmable calculators are permitted for this examination,—please indicate the make and model of your calculator on each answer book used.

1. (25 points) For each of the following equations, state whether *all* solutions are bounded for $t > 0$ and whether *all* solutions satisfy

$$\lim_{t \rightarrow +\infty} x(t) = 0.$$

While you needn't solve the equations explicitly, you should indicate how your answers were obtained.

(a) (5 points)

$$x''(t) + 2x'(t) + 3x(t) = 0$$

(b) (5 points)

$$x''(t) - 2x'(t) + 3x(t) = 0$$

(c) (5 points)

$$x''(t) + 4x(t) = 0$$

(d) (5 points)

$$x''(t) - 4x(t) = 0$$

(e) (5 points)

$$x'''(t) + x''(t) = 0$$

2. (25 points)

- (a) (5 points) How is the matrix exponential defined? *Note: I'm not asking how it's computed, I'm asking how it's defined.*
- (b) (5 points) What useful property of the ordinary exponential function does the matrix exponential fail to satisfy?
- (c) (15 points) Given that

$$\begin{pmatrix} -43 & -63 \\ 30 & 44 \end{pmatrix} = \begin{pmatrix} -7 & 3 \\ 5 & -2 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 2 & 3 \\ 5 & 7 \end{pmatrix},$$

express the general solution to the system

$$x'(t) = -43x(t) - 63y(t)$$

$$y'(t) = 30x(t) + 44y(t)$$

in terms of the initial values $x(0)$ and $y(0)$.

3. (25 points)

- (a) (5 points) Prove that

$$A = x^2 + y^2$$

and

$$B = x^2 + z^2$$

are invariants of the nonlinear system

$$x'(t) = y(t)z(t)$$

$$y'(t) = -x(t)z(t)$$

$$z'(t) = -x(t)y(t)$$

- (b) (10 points) Prove that all solutions of the system are bounded.
- (c) (10 points) Prove that $(0, 0, 0)$ is a stable equilibrium.

4. (25 points) Given that

$$x_1(t) = t^2 + 1$$

is a solution of the equation

$$(t^2 - 2t - 1)x''(t) + (-2t + 2)x'(t) + 2x(t) = 0,$$

find *all* solutions.

5. (25 points)

(a) (16 points) Define the following terms:

- i. Autonomous System
- ii. Equilibrium
- iii. Stable
- iv. Asymptotically stable
- v. Strictly stable
- vi. Linearisation
- vii. Lyapunov function
- viii. Strict Lyapunov function

(b) (9 points) Prove that

$$V(x, y) = x^2 + y^2$$

is a strict Lyapunov function for the equilibrium $(0, 0)$ of the system

$$x'(t) = -x(t) + y(t)$$

$$y'(t) = -x(t) - y(t)$$

6. (25 points) Chebyshev's equation is

$$(1 - t^2)x''(t) - t'x(t) + \nu^2 x(t) = 0.$$

For which values of ν does this equation have polynomial solutions? What are these solutions?