## MA1S12 (Timoney) Tutorial sheet 6b [March 3–7, 2014]

## Name: Solutions

1. Find the eigenvalues for the matrix

$$A = \begin{bmatrix} 7 & 0 & 0 \\ 0 & 5 & 1 \\ 0 & 1 & 5 \end{bmatrix}$$

Solution: The eigenvalues are the solutions of  $det(A - \lambda I_3) = 0$ . We compute

$$A - \lambda I_3 = \begin{bmatrix} 7 - \lambda & 0 & 0 \\ 0 & 5 - \lambda & 1 \\ 0 & 1 & 5 - \lambda \end{bmatrix}$$

and if we expand the determinant along the 1st row (or 1st column) we get

$$\det(A - \lambda I_3) = (7 - \lambda) \det \begin{bmatrix} 5 - \lambda & 1\\ 1 & 5 - \lambda \end{bmatrix} = (7 - \lambda)((5 - \lambda)^2 - 1)$$

The roots satisfy  $\lambda = 7$  and  $\lambda - 5 = \pm 1$ . So they are

7, 6 and 4.

Those are the eigenvalues.

2. For the same A, find unit eigenvectors (*i.e.* length 1 eigenvectors) for each eigenvalue.

Solution: For  $\lambda = 7$  we should solve the system that gives rise to doing Gaussian elimination on  $[A - 3I_3 : 0]$  or

$$\begin{bmatrix} 0 & 0 & 0 & : & 0 \\ 0 & -2 & 1 & : & 0 \\ 0 & 1 & -2 & : & 0 \end{bmatrix}$$

The eigenvector will turn out as

$$\begin{bmatrix} 1\\0\\0\end{bmatrix}$$

For  $\lambda = 6$  we should solve the system that gives rise to doing Gaussian elimination on  $[A - 6I_3:0]$  or

$$\begin{bmatrix} 1 & 0 & 0 & : & 0 \\ 0 & -1 & 1 & : & 0 \\ 0 & 1 & -1 & : & 0 \end{bmatrix}$$

An eigenvector will turn out as



For  $\lambda = 4$  we should solve the system that gives rise to doing Gaussian elimination on  $[A - 4I_3:0]$  or

$$\begin{bmatrix} 3 & 0 & 0 & : & 0 \\ 0 & 1 & 1 & : & 0 \\ 0 & 1 & 1 & : & 0 \end{bmatrix}$$

0

1 1

An eigenvector will turn out as

and we should normalise that to

$$\begin{bmatrix} 0\\ 1/\sqrt{2}\\ -1/\sqrt{2} \end{bmatrix}$$

3. For the same A, find an orthogonal matrix P and a diagonal matrix D so that A = PDP<sup>t</sup>.
Solution:

$$D = \begin{bmatrix} 7 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 4 \end{bmatrix}$$
$$P = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix}$$

(Other possible answers multiply the columns of P by  $\pm 1$  and you can permute 3, 6 and 4 on the diagonal of D if you apply the same permutation to the columns of P.

Richard M. Timoney

