MA 419 Assignment 2

Due 15 November 2006

1. Compute $f \star g$ where f and g are defined by

$$f(x) = \exp(-|x|)$$

and

$$g(x) = \begin{cases} 0 & \text{if } x < -1 \\ 1 & \text{if } -1 \le x \le 1 \\ 0 & \text{if } 1 < x \end{cases} .$$

2. The solution of the initial value problem for the homogeneous Wave Equation

$$u_{tt} - c^2 u_{xx} = 0$$
 $u|_{t=0} = \varphi$ $u_t|_{t=0} = \psi$

in the special case $\varphi = 0$ can be written in the form

$$u(t,\cdot) = K(t,\cdot) \star \psi$$

for an appropriate function K. What is this K?

3. A function f is said to be α -Hölder continuous if, for every x, there is an L and a $\delta > 0$ such that

$$|f(y) - f(x)| < L|y - x|^{\alpha}$$

whenever $|y - x| < \delta$. Show that if there is a positive α such that f is α -Hölder continuous then f is continuous. Show that if $f' \in L^p(\mathbf{R})$ for some $1 \leq p \leq \infty$ then f is α -Hölder continuous for

$$\alpha = 1 - \frac{1}{p}$$

4. Solve the Diffusion Equation

 $u_t - k u_{xx}$

with initial data

$$u(0,x) = e^{\lambda x}.$$

5. With S defined, as in lecture, by

$$S(t,x) = (4\pi kt)^{-1/2} \exp(-\frac{x^2}{4kt}),$$

prove that for all positive s and t

$$S(s+t,\cdot) = S(s,\cdot) \star S(t,\cdot).$$