

MA 216  
Assignment 2  
Due 7 November 2007

1. Rewrite each of the following higher order scalar equations as a first order system.

(a) Hook's law

$$mx''(t) + kx(t) = 0$$

(b) Bessel's equation

$$t^2 x''(t) + tx'(t) + (t^2 - \nu^2)x(t) = 0,$$

(c) The Laguerre Differential Equation

$$tx''(t) + (1 - t)x'(t) + \nu x(t) = 0.$$

2. It was shown in lecture that if  $AB = BA$  then each of the following equations is satisfied for all real  $t$ :

$$\exp(tA + tB) = \exp(tA) \exp(tB),$$

$$\exp(tA + tB) = \exp(tB) \exp(tA),$$

$$\exp(tA) \exp(tB) = \exp(tB) \exp(tA).$$

Prove that, conversely, if any of these three equations is satisfied for all  $t$  then  $AB = BA$ .

*Hint:* Differentiate twice.

3. Compute the following matrix exponentials:

(a)

$$\exp\left(t\begin{pmatrix} 13 & -5 \\ 30 & -12 \end{pmatrix}\right)$$

(b)

$$\exp\left(t\begin{pmatrix} 1 & 1 \\ -4 & 5 \end{pmatrix}\right)$$

(c)

$$\exp\left(t\begin{pmatrix} -2 & 2 \\ -13 & 8 \end{pmatrix}\right)$$

4. Find the solution to the initial value problem

$$x(0) = \xi_1 \quad x'(0) = \xi_2$$

for the general linear constant coefficient second order scalar equation

$$c_2 x''(t) + c_1 x'(t) + c_0 x(t) = 0$$

by reducing it to a first order system.