

MA 4448
Assignment 4
Due 31 March 2011

Id: 4448-1011-4.m4,v 1.6 2011/04/26 13:05:01 john Exp john

1. The Kerr metric is, in coordinates t, r, θ, φ ,

$$\begin{aligned}g_{tt} &= -c^2 \left(1 - \frac{2G\bar{m}r}{c^2 r^2 \left(1 + \frac{j^2 \cos^2 \theta}{\bar{m}^2 c^2 r^2} \right)} \right) \\g_{t\varphi} = g_{\varphi t} &= -\frac{2Gj \sin^2 \theta}{c^2 r \left(1 + \frac{j^2 \cos^2 \theta}{\bar{m}^2 c^2 r^2} \right)} \\g_{rr} &= \frac{1 + \frac{j^2 \cos^2 \theta}{c^2 \bar{m}^2 r^2}}{1 - \frac{2G\bar{m}}{c^2 r} + \frac{j^2}{c^2 \bar{m}^2 r^2}} \\g_{\theta\theta} &= r^2 \left(1 + \frac{j^2 \cos^2 \theta}{\bar{m}^2 c^2 r^2} \right) \\g_{\varphi\varphi} &= \left(r^2 + \frac{j^2}{c^2 \bar{m}^2} + \frac{2Gj^2 \sin^2 \theta}{c^4 \bar{m} r \left(1 + \frac{j^2 \cos^2 \theta}{\bar{m}^2 c^2 r^2} \right)} \right) \sin^2 \theta.\end{aligned}$$

Here J is the angular momentum of the central mass and \bar{m} is its mass. G is the Newtonian gravitational constant. All other entries of the metric tensor are zero. Find the redshift between an arbitrary pair of stationary observers.

Note: Don't be frightened by the hideousness of the metric. The problem is in fact really easy.

2. For the Kerr metric, as for the Reissner-Nordström, $\theta \rightarrow \pi - \theta$ is a symmetry, so if $\theta = \pi/2$ initially and $\dot{\theta} = 0$ then $\theta = \pi/2$ for all time. We *don't* have full rotational symmetry so we can *not* say that $\theta = \pi/2$ without loss of generality. But the case $\theta = \pi/2$ is still physically interesting. This problem is equivalent to studying the motion of a particle in a three dimensional space time with coordinates t, r and φ with metric tensor obtained by setting $\theta = \pi/2$ in the equations above and then dropping $g_{\theta\theta}$:

$$g_{tt} = -c^2 \left(1 - \frac{2G\bar{m}}{c^2 r} \right)$$

$$g_{t\varphi} = g_{\varphi t} = -\frac{2Gj}{c^2 r}$$

$$g_{rr} = \frac{1}{1 - \frac{2G\bar{m}}{c^2 r} + \frac{j^2}{c^2 \bar{m}^2 r^2}}$$

$$g_{\varphi\varphi} = \left(r^2 + \frac{j^2}{c^2 \bar{m}^2} + \frac{2Gj^2}{c^4 \bar{m} r} \right).$$

What is the inverse metric?

3. What are the canonical momenta and Hamiltonian for an uncharged test particle of mass m on a spacetime with metric as in the previous problem? What algebraic relation do they satisfy?