# MA 4448
# Assignment 2
# Due 23 February 2011

`Id: 4448-1011-2.m4,v 1.2 2011/03/03 11:49:43 john Exp john`

1. With spherical coordinates $(t, r, \theta, \varphi)$, let $g_{(0)} = \partial/\partial t$, $g_{(1)} = \partial/\partial r$, $g_{(2)} = r^{-1}\partial/\partial\theta$, and $g_{(3)} = (r \sin\theta)^{-1}\partial/\partial\varphi$.

   (a) Show that $\{g_{(0)}, g_{(1)}, g_{(2)}, g_{(3)}\}$ is a local frame.
   *Solution:* We know that $\{\partial/\partial t, \partial/\partial x, \partial/\partial y, \partial/\partial z\}$ is a basis at each point. $t, r, \theta, \varphi$ are related to $t, x, y, z$ by

   $$t = t \quad x = r\sin\theta\cos\varphi \quad y = r\sin\theta\sin\varphi \quad z = r\cos\theta$$

   By the chain rule,

   $$\begin{aligned}
   \begin{pmatrix} \frac{\partial}{\partial t} \\ \frac{\partial}{\partial r} \\ \frac{\partial}{\partial \theta} \\ \frac{\partial}{\partial \varphi} \end{pmatrix} &= \begin{pmatrix} \frac{\partial t}{\partial t} & \frac{\partial x}{\partial t} & \frac{\partial y}{\partial t} & \frac{\partial z}{\partial t} \\ \frac{\partial t}{\partial r} & \frac{\partial x}{\partial r} & \frac{\partial y}{\partial r} & \frac{\partial z}{\partial r} \\ \frac{\partial t}{\partial \theta} & \frac{\partial x}{\partial \theta} & \frac{\partial y}{\partial \theta} & \frac{\partial z}{\partial \theta} \\ \frac{\partial t}{\partial \varphi} & \frac{\partial x}{\partial \varphi} & \frac{\partial y}{\partial \varphi} & \frac{\partial z}{\partial \varphi} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial t} \\ \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \end{pmatrix} \\
   &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \sin\theta\cos\varphi & \sin\theta\sin\varphi & \cos\theta \\ 0 & r\cos\theta\cos\varphi & r\cos\theta\sin\varphi & -r\sin\varphi \\ 0 & -r\sin\theta\sin\varphi & r\sin\theta\cos\varphi & 0 \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial t} \\ \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \end{pmatrix}.
   \end{aligned}$$

   Then

   $$\begin{aligned}
   \begin{pmatrix} g_{(0)} \\ g_{(1)} \\ g_{(2)} \\ g_{(3)} \end{pmatrix} &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & r^{-1} & 0 \\ 0 & 0 & 0 & (r\sin\theta)^{-1} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial t} \\ \frac{\partial}{\partial r} \\ \frac{\partial}{\partial \theta} \\ \frac{\partial}{\partial \varphi} \end{pmatrix} \\
   &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \sin\theta\cos\varphi & \sin\theta\sin\varphi & \cos\theta \\ 0 & \cos\theta\cos\varphi & \cos\theta\sin\varphi & -\sin\varphi \\ 0 & \sin\varphi & \cos\varphi & 0 \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial t} \\ \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \end{pmatrix}.
   \end{aligned}$$

The matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \sin\theta\cos\varphi & \sin\theta\sin\varphi & \cos\theta \\ 0 & \cos\theta\cos\varphi & \cos\theta\sin\varphi & -\sin\varphi \\ 0 & \sin\varphi & \cos\varphi & 0 \end{pmatrix}$$

has determinant 1, and hence is invertible, so $\{g_{(0)}, g_{(1)}, g_{(2)}, g_{(3)}\}$ is a basis because $\{\partial/\partial t, \partial/\partial r, \partial/\partial\theta, \partial/\partial\varphi\}$ is.

(b) Compute the metric coefficients in this frame, and show that this frame is a tetrad.

*Solution:* Let $\cdot$ denote inner products. Then

$$g_{(0)} \cdot g_{(0)} = \frac{\partial}{\partial t} \cdot \frac{\partial}{\partial t} = g_{tt} = -c^2,$$

$$g_{(1)} \cdot g_{(1)} = \frac{\partial}{\partial r} \cdot \frac{\partial}{\partial r} = g_{rr} = 1,$$

$$g_{(2)} \cdot g_{(2)} = \frac{1}{r}\frac{\partial}{\partial\theta} \cdot \frac{1}{r}\frac{\partial}{\partial\theta} = \frac{1}{r^2} g_{\theta\theta} = 1,$$

and

$$g_{(3)} \cdot g_{(3)} = \frac{1}{r\sin\theta} l \frac{\partial}{\partial\varphi} \cdot \frac{1}{r\sin\theta}\frac{\partial}{\partial\varphi} = \frac{1}{r^2\sin^2\theta} g_{\varphi\varphi} = 1.$$

The non-diagonal commponents are all zero. Since the components of the metric are all constant in this frame it is, by definition, a tetrad.

(c) Compute the commutation coefficients, defined by

$$[g_{(b)}, g_{(c)}] = \gamma^a_{bc} g_{(a)}.$$

*Solution:*

$$\begin{aligned} \left[g_{(2)}, g_{(3)}\right] = -\left[g_{(2)}, g_{(3)}\right] &= \left[\frac{1}{r}\frac{\partial}{\partial\theta}, \frac{1}{r\sin\theta}\frac{\partial}{\partial\varphi}\right] \\ &= \frac{1}{r}\frac{\partial}{\partial\theta}\left(\frac{1}{r\sin\theta}\frac{\partial}{\partial\varphi}\right) - \frac{1}{r\sin\theta}\frac{\partial}{\partial\varphi}\left(\frac{1}{r}\frac{\partial}{\partial\theta}\right) \\ &= \frac{\cos\theta}{r^2\sin^2\theta}\frac{\partial}{\partial\varphi} = r^{-1}\cot\theta g_{(2)} \end{aligned}$$

so

$$\gamma^2_{23} = -\gamma^2_{32} = r^{-1}\cot\theta$$

and

$$\gamma^a_{23} = -\gamma^a_{32} = 0$$

if $a \neq 2$. The other commutators are similar, but simpler. The other non-zero components are

$$\gamma^2_{12} = -\gamma^2_{21} = -r^{-1} = \gamma^3_{13} = -\gamma^3_{31}.$$

2. Suppose $X = X^a\partial/\partial x^a$ and $Y = Y^b\partial/\partial x^b$ are vector fields and $\alpha = \alpha_c dx^c$ is a 1-form. Prove that

$$(d\alpha)(X, Y) = X\alpha(Y) - Y\alpha(X) - \alpha([X, Y]).$$

*Solution:*

By definition

$$(d\alpha)(X, Y) = (d\alpha)_{ab}X^aY^b = \epsilon^{cd}ab\frac{\partial}{\partial x^c}\alpha_d X^aY^b = (\alpha_{b,a} - \alpha_{a,b})X^aY^b.$$

Now

$$\alpha(X) = \alpha_c dx^c(X^a\partial/\partial x^a) = \alpha_c X^a dx^c(\partial/\partial x^a) = \alpha_c X^a g^c_a = \alpha_a X^a.$$

Similarly,

$$\alpha(Y) = \alpha_b Y^b.$$

Then

$$X\alpha(Y) = X^a\partial/\partial x^a(\alpha_b Y^b) = X^a(\alpha_{b,a}Y^b + \alpha_b Y^b_{,a}) = \alpha_{b,a}X^aY^b + \alpha_b X^aY^b_{,a}$$

and

$$Y\alpha(X) = \alpha_{a,b}X^aY^b + \alpha_a X^a_{,b}Y^b.$$

By definition,

$$[X, Y] = XY - YX$$

$$XY = X^a\partial/\partial x^a(Y^b\partial/\partial x^b) + X^aY^b\partial^2/\partial x^a\partial x^b$$

and

$$YX = Y^b\partial/\partial x^b(X^a\partial/\partial x^a) + X^aY^b\partial^2/\partial x^b\partial x^a$$

so

$$[X, Y] = X^aY^b_{,a}\partial/\partial x^b - X^a_{,b}Y^b\partial/\partial x^a$$

and

$$\alpha([X, Y]) = \alpha_c X^aY^b_{,a}dx^c(\partial/\partial x^b) - \alpha_c X^a_{,b}Y^b dx^c(\partial/\partial x^a) = \alpha_b X^aY^b_{,a} - \alpha_a X^a_{,b}Y^b.$$

Combining the results above

$$X\alpha(Y) - Y\alpha(X) - \alpha([X, Y]) = (\alpha_{b,a} - \alpha_{a,b})X^aY^b.$$

This is the same as our expression for $(d\alpha)(X, Y)$.

3. Suppose $\{g_{(0)}, g_{(1)}, g_{(2)}, g_{(3)}\}$ is a general frame, *i.e.* that it's values at $x$ are a basis for $T_x\mathcal{M}$. Let $\{g^{(0)}, g^{(1)}, g^{(2)}, g^{(3)}\}$ be the dual frame, *i.e.* its values at $x$ are the dual basis for $T_x^*\mathcal{M} = \Omega_x^1\mathcal{M}$. Then $\{g^{(0)} \wedge g^{(1)} = -g^{(1)} \wedge g^{(0)}, g^{(0)} \wedge g^{(2)} = -g^{(2)} \wedge g^{(0)}, g^{(0)} \wedge g^{(3)} = -g^{(3)} \wedge g^{(0)}, g^{(1)} \wedge g^{(2)} = -g^{(2)} \wedge g^{(1)}, g^{(1)} \wedge g^{(3)} = -g^{(3)} \wedge g^{(1)}, g^{(2)} \wedge g^{(3)} = -g^{(3)} \wedge g^{(2)}\}$ is, at each point $x$, a basis for $\Omega_x^2\mathcal{M}$. What are $\{dg^{(0)}, dg^{(1)}, dg^{(2)}, dg^{(3)}\}$ in this basis?
*Note:* You may use the result of the previous problem even if you didn't succeed in proving it.
*Solution:*

Apply the result of the previous problem to $X = g_{(a)}$, $Y = g_{(b)}$ and $\alpha = g^{(c)}$.

$$(dg^{(c)})(g_{(a)}, g_{(b)}) = g_{(a)}(g^{(c)}(g_{(b)})) - g_{(b)}(g^{(c)}(g_{(a)})) - g^{(c)}([g_{(a)}, g_{(b)}]).$$

Now

$$g^{(c)}(g_{(a)}) = g_{(a)}^{(c)},$$

which is constant, and hence has zero derivatives. Thus

$$g_{(a)}(g^{(c)}(g_{(b)})) = 0$$

Similarly,

$$g_{(b)}(g^{(c)}(g_{(a)})) = 0$$

By definition,

$$[g_{(a)}, g_{(b)}] = \gamma_{ab}^d g_{(d)}$$

and hence

$$g^{(c)}([g_{(a)}, g_{(b)}]) = \gamma_{ab}^d g^{(c)}(g_{(d)}) = \gamma_{ab}^d g_{(d)}^{(c)} = \gamma_{ab}^c.$$

Therefore

$$(dg^{(c)})(g_{(a)}, g_{(b)}) = -\gamma_{ab}^c$$

and

$$dg^{(c)} = -\gamma_{ab}^c g^{(a)} \wedge g^{(b)}.$$