

MA 3421  
Assignment 5  
Due 21 November 2012

Id: 3421-f2012-5.m4,v 1.1 2012/11/15 14:27:14 john Exp john

1. It was shown in lecture that if  $\sum_{k,l} |\alpha_{k,l}|^2 < \infty$  then the linear transformation  $A: \ell^2 \rightarrow \ell^2$  defined by  $(A\xi)_k = \sum_l \alpha_{k,l} \xi_l$  is compact. The condition  $\sum_{k,l} |\alpha_{k,l}|^2 < \infty$  is sufficient for compactness, but not necessary. To show this, consider the operator  $A$  with

$$\alpha_{k,l} = \begin{cases} k^{-s} & \text{if } k = l, \\ 0 & \text{if } k \neq l, \end{cases}$$

where  $s \geq 0$ .

- (a) For which values of  $s$  is  $A$  bounded?
  - (b) For which values of  $s$  is  $A$  compact?
  - (c) For which values of  $s$  is  $A$  symmetric?
  - (d) For which values of  $s$  is  $\sum_{k,l} |\alpha_{k,l}|^2 < \infty$ ?
2. Let  $L$  and  $R$  be the left and right unilateral shift operators on  $\ell^2$ ,

$$(L\xi)_n = \xi_{n+1} \quad (R\xi)_n = \begin{cases} 0 & \text{if } n = 1, \\ \xi_{n-1} & \text{if } n > 1. \end{cases}$$

- (a) Is  $L$  compact?

*Id: 3421-f2012-5.m4, v 1.1 2012/11/15 14:27:14 john Exp john 2*

(b) Is  $R$  compact?

(c) Is  $L$  symmetric?

(d) Is  $R$  symmetric?

(e) What are the eigenvalues and eigenvectors of  $L$ ?

(f) What are the eigenvalues and eigenvectors of  $R$ ?