

MA 2326
Assignment 5
Due 20 March 2014

Id: 2326-s2014-5.m4,v 1.1 2014/03/13 15:50:18 john Exp john

1. Find the equilibria of the autonomous system

$$x'(t) = x(x^3 - 2y^3), \quad y' = y(2x^3 - y^3).$$

2. Suppose that U and V are open subsets of \mathbf{R}^m and that $\varphi: U \rightarrow V$ and $\psi: V \rightarrow U$ are continuously differentiable and that ψ is the inverse of φ . Suppose also that $F: U \rightarrow \mathbf{R}^m$ and that $G: V \rightarrow \mathbf{R}^m$ is defined by

$$G_j(y) = \sum_{k=1}^m \frac{\partial \varphi_j}{\partial x_k}(\psi(y)) F_k(\psi(y)).$$

- (a) Prove that $x: I \rightarrow U$ is a solution to the autonomous system

$$x'(t) = F(x(t))$$

if and only if $y: I \rightarrow V$, defined by

$$y(t) = \varphi(x(t)),$$

is a solution to the autonomous system

$$y'(t) = G(y(t)).$$

I is an interval.

- (b) Prove that ξ is an equilibrium of $x'(t) = F(x(t))$ if and only if $\eta = \varphi(\xi)$ is an equilibrium of $y'(t) = G(y(t))$.
- (c) Prove that ξ is a stable equilibrium if and only if η is.
- (d) Prove that ξ is a strictly stable equilibrium if and only if η is.