#### Course 2318 2011

### Sheet 1

Due: at the end of the tutorial on Thursday of the next week

## Exercise 1

Which of the following sets can always be transformed into each other by an euclidean transformation?

- (i) Two lines in the plane.
- (ii) Two pairs of intersecting lines in the plane.
- (iii) Two pairs of intersecting lines in the plane forming the same angles. Justify your answer.

### Exercise 2

Determine the type (in the classification: ellipse, hyperbola etc.) of the following quadric:

- (i) xy + x 1 = 0;
- (ii)  $x^2 2xy + y^2 + x 1 = 0;$
- (iii)  $x^2 2xy + y^2 1 = 0;$
- (iv)  $x^2 + y^2 x y + 1 = 0;$

# Exercise 3

In  $\mathbb{R}^3$  consider two planes A and B given by Z = 1 and X = 1 respectively. Consider inhomogeneous coordinates (x, y) = (X/Z, Y/Z) on A and (y', z') = (Y/X, Z/X) on B. Let  $L \subset A$  be the line given by y = ax + b.

- (i) Describe by a homogeneous equation the projective line  $\tilde{L}$  in  $\mathbb{P}^2$  induced by L, i.e. containing all the lines passing through 0 and intersecting L.
- (ii) Give an equation for the line induced by  $\widetilde{L}$  on the plane B using the coordinates (y', z').
- (iii) What is the "infinity point" of  $\tilde{L}$ , i.e. the point in  $\mathbb{P}^2$  which does not correspond to any point of L?
- (iv) For which a and b, the latter "infinity point" corresponds to a point of B?